

Drive[®] XL

Herbicide

For unsurpassed post emergent control of Summer grass, White clover and suppression of Kikuyu in turf.

General Description

Drive[®] XL herbicide powered by Quinclorac, is the latest technology in providing unsurpassed control of Summer grass, White clover and suppression of Kikuyu in turf grasses. A proprietary, water based formulation, **Drive XL** provides faster more effective weed control, delivers quicker plant uptake and is rainfast in as little as one hour.

Features

Benefits

Water based formulation

Provides faster more effective control through quicker plant uptake of the active ingredient.

Unsurpassed performance

Drive XL provides superior control of Summer and White clover and suppression of Kikuyu.

Rainfast and fast drying

Drive XL is rainfast within one hour of application.

Suitable for use on cool and warm grass species

Drive XL is registered for use on a range of established turf grass species – refer to product label.

Application Information

Apply **Drive XL** to actively growing weeds. Use **Drive XL** at a rate of 4.6L per 400L tank mix. 400L of tank mix is sufficient to cover one hectare. Avoid overlaps during application, which can be achieved through the use of a spray marker dye such as **Turf Mark[®] Blue**. For spot spray application, apply to point of run off. Separate applications should be made if all target weeds are not at the correct growth stage for treatment at the same time. It is recommended not to mow 2 days before or after applying **Drive XL** to maximise weed control and minimise turf injury. Mix with a high quality Methylated Seed Oil (MSO) e.g. Hasten[®] at 1% concentration (v/v) for superior performance.

Packaging

Available in 5L packs

Product Update

Toll Free: 1800 558 399

agro.basf.com.au

 **BASF**
We create chemistry

Drive[®] XL

Herbicide

For unsurpassed post emergent control of summer grass, white clover and suppression of kikuyu in turf

SITUATION	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
<p>Established turf of: Common couch (<i>Cynodon dactylon</i>), hybrid couch (<i>Cynodon dactylon</i> x <i>C. transvaalensis</i>), Japanese lawngress (<i>Zoysia japonica</i>), marine couch (<i>Sporobolus virginicus</i>), perennial ryegrass (<i>Lolium perenne</i>) creeping bentgrass (<i>Agrostis stolonifera</i>), Kentucky Bluegrass (<i>Poa pratensis</i>)</p> <p>See also “Protection of crops, native and other non- target plants” section</p>	<p>Summer grass (<i>Digitaria</i> spp.)</p>	<p>4.6L PER 400L WATER</p>	<p><u>One application:</u> Weeds at cotyledon stage up to prior to second tiller OR Weeds at 5 tillers and greater</p> <p><u>Two applications, 21 to 28 days apart:</u> Weeds from 2 to 4 tillers The addition of a crop oil adjuvant may improve weed control in unfavourable weather conditions.</p>
	<p>Kikuyu (<i>Pennisetum clandestinum</i>) (Suppression only)</p>		<p>For Kikuyu suppression, apply twice, 14 to 21 days apart. Best results are gained from Autumn applications. The addition of a crop oil adjuvant may improve weed control in unfavourable weather conditions.</p>
	<p>White clover (<i>Trifolium repens</i>)</p>		<p>Apply to actively growing weeds.</p>

ALWAYS READ AND FOLLOW LABEL DIRECTIONS.

This advertisement is intended as general advice. Disclaimer: The information submitted in this publication is based on current BASF knowledge and experience. In view of the many factors that may affect its application, this data does not relieve the user from carrying out their own tests. The data does not imply assurance of certain properties or of suitability for a specific purpose. It is the responsibility of the user to ensure that any proprietary rights and existing laws and legislation are observed.

©Copyright BASF 2015 ©Registered trademark of BASF

BASF Australia Limited, Level 12, 28 Freshwater Place, Southbank Victoria 3006, Australia.

Toll Free: 1800 558 399

agro.basf.com.au

 **BASF**
We create chemistry