

CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FARMALINX

Scanner 500 SC

HERBICIDE

ACTIVE CONSTITUENT: 500 g/L ETHOFUMESATE

GROUP **J** HERBICIDE

For selective weed control in beet crops, certain non-crop areas, onions, oilseed poppies, ryegrass pastures and seed crops and established turf as specified in the Directions for Use table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT.

farmalinx

FARMALINX Pty Ltd ABN 95 134 353 245
Level 25, Suite 2506, Tower 2, 101 Grafton Street, Bondi Junction NSW 2022
T: 02 9389 2455 F: 02 9389 2844 W: www.farmalinx.com

CONTENTS

5 L

FARMALINX

Scanner 500 SC

HERBICIDE

UN No 3082

ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S.

STORAGE AND DISPOSAL: Store in the closed original container in a dry, well-ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight. It is recommended that the storage temperature never exceed 30°C. Triple or (preferably) pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycle or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulation. Do not burn empty containers or product.

SAFETY DIRECTIONS: May irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use.

FIRST AID: If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126.

MATERIAL SAFETY DATA SHEET: Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FARMALINX SCANNER 500 SC Herbicide is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 9389 2455.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to the label instructions, or under off-label permits not endorsed by Farmalinx Pty Ltd or under abnormal conditions. Farmalinx Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

APVMA Approval No.: 69025/59668



HAZCHEM CODE: 3Z
PACKAGING GROUP: III



**IN A TRANSPORT
EMERGENCY
DIAL 000**
POLICE OR FIRE BRIGADE

SPECIALIST ADVICE IN AN EMERGENCY ONLY

1800 024 973

ALL HOURS – AUSTRALIA WIDE



Batch No.

Date of Manufacture

farmalinx

FARMALINX

Scanner 500 SC

HERBICIDE

DIRECTIONS FOR USE

RESTRAINTS:

DO NOT apply if rain is imminent or falling.
DO NOT rotate with crops other than beet crops for 6 months after application.

CROP OR SITUATION	WEEDS CONTROLLED	STATE	RATE	WHP	CRITICAL COMMENTS
Beet crops eg., beetroot, fodder beet, mangolds, sugar beet	Barley Grass, Chickweed, Cleavers, Fat-hen, Fumitory, Prairie Grass, Redshank (<i>Amaranthus</i> spp.), Shepherd's Purse, Summer Grass, Winter Grass, Wireweed	Qld, NSW, ACT, Vic, Tas, SA only	4 or 6 L/ha	-	<p>Pre-emergence: Apply before weeds or crop emerge, either pre- or post-sowing of crops. Use high rate on heavy soil or where <i>Amaranthus</i> spp., Prairie Grass or Wireweed are a problem. DO NOT re-treat area at these rates.</p> <p>Post-emergence: Crop Stage: Apply at 2 to 4 leaf stage. Weed Stage: Apply when weeds are no larger than 2-4 leaf stage, either as a band spray or overall application. DO NOT spray beets using high pressure. DO NOT spray beet crops when foliage is wet.</p>
			2 L plus 5 L Betanal*/ha		
Oilseed poppy	Barley Grass, certain broadleaved and grass weeds particularly Clovers, Cruciferous Weeds, Fat-hen, Fumitory, Winter Grass, Wireweed	Tas only	400 mL to 2 L/ha		The range of control is extended with spray programs in combinations with asulam, diclofop-methyl and diquat as recommended by poppy contracting companies or the Department of Primary Industry. DO NOT add wetting agent.
Onions	Barley Grass, certain broadleaved and grass weeds, Clovers, particularly Cruciferous Weeds, Fat-hen, Fumitory, Winter Grass, Wireweed	Tas only	600 mL to 1.2 L/ha	13 weeks	Apply after 2 leaf stage of crop. DO NOT add wetting agent. Follow up with spray programs using alternative herbicides as recommended by the Department of Primary Industry, or contracting companies for use in onions.
			1 L to 1.2 L plus 2.8 kg active constituent chlorpropham/ha		This mixture, and follow-up spray programs with alternative herbicides should be used according to recommendations by the Department of Primary Industry or contracting companies. Apply after 2 leaf stage of crop. DO NOT add wetting agent to SCANNER 500 SC.

CROP OR SITUATION	WEEDS CONTROLLED	STATE	RATE	WHP	CRITICAL COMMENTS	
Ryegrass pasture and seed crops also, Non-Crop areas (stock camps, tree plantations, fence lines)	Barley Grass, Cleavers, emerged Brome Grass, seedling Chickweed, Shepherd's Purse, Storksbill, Winter Grass	Vic, Tas only	3 L/ha	7 days	Apply (April-July) to moist soil after main germination of Barley Grass. DO NOT apply once Barley Grass reaches green flowerhead stage. DO NOT treat pastures where clover production is important. Use this rate when: 1. Annual rainfall is 250-750 mm. 2. Barley Grass is at very early growth stage (April to May). 3. Pasture has been closely grazed (less than 5 cm high). 4. Soil is low in organic matter (less than 10%). 5. There is a high proportion of ryegrass in pasture	
			4 L/ha			Use this rate when: 1. Annual rainfall exceeds 750 mm. 2. Barley grass is at early vegetative growth stage (April-July). 3. Pasture has been closely grazed (5-10 cm high). 4. Dung accumulations are present.
			300 mL per 100 L water			Spot Treatment: Apply by handgun or knapsack sprayer. Thoroughly wet vegetation.
Established turf; bent grasses (including brown top bent), Kentucky blue-grass, kikuyu, Queensland blue couch, ryegrass	Winter Grass (<i>Poa annua</i>)	NSW, ACT, Vic, Tas, SA, WA only	800 mL in 400 to 600 L water/ha OR 8 mL in 4 to 6 L water per 100 m ²	DO NOT GRAZE OR FEED TO ANIMALS	Apply first application one month before main Autumn germination (ie. apply in February to April but up to May in northern NSW). Repeat at monthly intervals for 3 to 4 months. In dry conditions, watering will improve residual action. After spraying, wait at least 12 hours before watering. Avoid flooding or ponding of treated turf. Preferably mow immediately before spraying, or leave as long as possible after spraying before next mowing (1 to 2 weeks if practicable). Water volume is not critical but avoid excessive overlapping and dripping from spray equipment onto turf. DO NOT add other herbicides or surfactants to this product for this use unless specifically recommended.	

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

www.farmalinx.com

CONTINUED OVERLEAF

STORAGE AND DISPOSAL Store in the closed original container in a dry, well-ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight. It is recommended that the storage temperature never exceed 30°C. Triple or (preferably) pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of unlabelled chemicals on-site. If recycling, replace cap and return clean containers to recycle or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If the empty packaging is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulation. Do not burn empty containers or product.

SAFETY DIRECTIONS May irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use.

FIRST AID If poisoning occurs contact a doctor or Poisons Information Centre. If Phone Australia 131 126.

MATERIAL SAFETY DATA SHEET Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FARMALINX SCANNER 500 SC Herbicide is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 9389 2455.

Other trademarks arising from incorrect storage, handling or use.

MIXING SCANNER 500 SC is a suspension concentrate which mixes easily with water. Prior to pouring, gently invert or roll container to ensure all active constituent is in suspension. Part-fill the spray tank with clean water. Add the correct amount of SCANNER 500 SC and top up with water to the required level. Thorough agitation of the mixture is necessary before spraying. Mix only enough chemical for immediate use. When using a mixture of SCANNER 500 SC and Betanal, half-fill spray vat with water then stir or agitate while adding the required quantities. Top up with water and stir again. Use as soon as possible after mixing. DO NOT add other herbicides or surfactants unless specifically recommended.

APPLICATION Apply in a suitable volume of water to give good boom spray. When used pre-emergence, apply to moist soil surface. Ensure that the nozzles are set at the correct height to give even coverage ie. 50-60 cm above the pasture or crop. Use a nozzle and pressure combination to give a fine spray (200-300 kPa). Make sure nozzle tips are clean, undamaged and all the correct type and size. Impact nozzles or broadcast nozzles clusters should NOT be used. DO NOT use filter gauze finer than 0.3 mm. After use, wash all equipment thoroughly with water containing liquid detergent. Spray out completely, then rinse twice with clean water. To allow air access, remove nozzles and open spray tank lid.

RESISTANT WEEDS WARNING FARMALINX SCANNER 500 SC is a member of the HERBICIDE J GROUP FUNGICIDE. SCANNER 500 SC has the Benzotriazin group of herbicides. SCANNER 500 SC is a Group J herbicide. Some naturally occurring weed biotypes resistant to SCANNER 500 SC and other Group J herbicides may exist through normal genetic variability in any weed population. The resistant individuals are used repeatedly. These resistant weeds will not be controlled by SCANNER 500 SC or other Group J herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, FARMALINX Pty Ltd accepts no liability for any losses that may result from the failure of SCANNER 500 SC to control resistant weeds. Ensure that the correct amount of SCANNER 500 SC is applied to detect and control resistant weeds. Do NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

WITHOLDING PERIODS (WHP):
ONIONS: DO NOT HARVEST FOR 13 WEEKS AFTER APPLICATION.
RYEGRASS PASTURE AND SEED CROPS:
DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.
TURF: DO NOT GRAZE TREATED AREAS OR FEED LAWN (TURF) CLIPPINGS FROM TREATED AREAS TO ANIMALS, INCLUDING POULTRY.

GENERAL INSTRUCTIONS
 The product is a pre- and post-emergence herbicide for control of certain broad-leaved weeds and grasses. For post-emergence applications, weeds should be at the seedling to 4 leaf stage. Prior to sowing any follow-up crops other than beet or ryegrass, those areas previously treated should be thoroughly tilled (including ploughing).
 1. BEET CROPS: Slight foliage distortion to beet may occur following treatment. These effects are short-lived and do not affect crop yield. Pre-emergence treatment with this product may be followed by post-emergence application of Betanal* or a mixture of both.
 2. USE OF SCANNER 500 SC IN MIXTURE WITH Betanal ON BEET CROPS: Before using this mixture carefully read the Betanal label; note warning about use in temperatures above 30°C. For best results the interval between treatment and subsequent rainfall or irrigation should be at least six (6) hours. In Tasmania, this treatment should follow the use of a pre-emergence herbicide. This mixture is recommended to provide a wider spectrum of post-emergence weed control and longer residual activity than Betanal used alone. Weeds controlled include additional weeds listed on the Betanal label.
 3. RYEGRASS: This product will reduce clover stands by 40-60%. For best control of barley grass, apply to closely grazed pasture (3-5 cm high). Raw organic matter can reduce efficacy and retreatment may be necessary. Remove organic litter and harrow large dung pats before application. In areas of dense Barley Grass, drill or oversow with ryegrass as soon as possible after treatment.
 4. TURF: This selective herbicide should be applied before, or early after germination or emergence of Winter Grass (*Poa annua*) when it occurs in fine turf situations. It is selective in ryegrass, fescues, Kentucky bluegrass, Queensland blue couch, kikuyu, bent grasses (including brown top bent) and couch, when applied as directed. Applications are made at monthly intervals for 3-4 months to control winter grass gradually, while allowing the desirable turf grasses to take over without unsightly bare patches of dead Winter Grass. Overseeding and reseeding should NOT take place within one month of spraying.

farmalinx

FARMALINX Pty Ltd ABN 95 134 353 245
 Level 25, Suite 2506, Tower 2,
 101 Grafton Street, Bondi Junction NSW 2022
 T: 02 9389 2455 F: 02 9389 2844
 Website: www.farmalinx.com

APVMA Approval No. 69025/59668

CAUTION
 KEEP OUT OF REACH OF CHILDREN
 READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Scanner 500 SC

HERBICIDE

ACTIVE CONSTITUENT: 500 g/L ETHOFUMESATE

GROUP J HERBICIDE

For selective weed control in beet crops, certain non-crop areas, onions, oilseed poppies, ryegrass pastures and seed crops and established turf as specified in the Directions for Use table.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

farmalinx

FARMALINX Pty Ltd ABN 95 134 353 245
 Level 25, Suite 2506, Tower 2, 101 Grafton Street,
 Bondi Junction NSW 2022