

DIRECTIONS FOR USE**Restraints****DO NOT APPLY if weeds or turf are under stress or when frosts are imminent****DO NOT USE if rain is expected within 4 hours.****DO NOT USE on bent grass or winter grass golf and bowling greens.**

Situation	Weeds Controlled	Rate	Critical Comments
Turf	Broadleaf weeds including; White Clover (<i>Trifolium repens</i>) Plantain (<i>Plantago lanceolata</i>) Capeweed (<i>Arctotheca calendula</i>) Cat's ear (<i>Hypochoeris radicata</i>) Bin-dii (Jo-Jo, (<i>Soliva sessilis</i>) Onehunga) Cudweed (<i>Gnaphalium</i> spp.) Creeping oxalis (<i>Oxalis corniculata</i>)	5 L in 200-500 L water/ha	Apply to actively growing weeds. Transient discolouration may occur on kikuyu and carpet grass and Queensland blue couch. Varietal differences in certain buffalo grasses (eg. ST85) may produce more pronounced effects and it is recommended that small areas be tested for turf safety before large-scale application occurs. Avoid exceeding application rates through overlapping sprayed areas. Use a surfactant for difficult to wet weeds.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**GENERAL INSTRUCTIONS**

For best results the turf and weeds should be actively growing at the time of treatment. Complete weed death may take up to 6 weeks. Faster results can be expected on smaller weeds. Due to residual effects of diflufenican, overseeding treated areas within 3 months of application may inhibit establishment of the turfgrass.

Mixing

Shake original container before use. Agitate continuously to ensure thorough mixing before and during application. Only mix sufficient chemical for each day's work. Surfactants (if used) should be added after Spearhead and water are mixed when the spray-tank is almost full.

Mowing

Avoid mowing during the 3 to 4 days preceding or following treatment.

Cleaning Spray Equipment

THOROUGHLY CLEAN ALL APPLICATION EQUIPMENT AFTER USE. Rinse water should be discharged onto a designated disposal area or if unavailable onto unused land away from desirable plants and water-courses.

Partial Cleaning (before spraying tolerant turf): After using Spearhead, empty the tank completely and drain the whole system. Thoroughly wash inside the tank with a pressure hose. Quarter fill the tank with clean water and circulate through the pump, line, hoses and nozzles. Drain and repeat procedure twice.

Complete cleaning (before spraying susceptible plants): After using Spearhead empty the tank completely and drain the whole system. Thoroughly wash inside the tank with a pressure hose. Quarter fill the tank with clean water and circulate through the pump, line, hoses and nozzles, then drain. Quarter fill the tank again and add an alkaline detergent at 500 mL/100 L water and circulate through

the system for at least 15 minutes. Drain, remove filters and nozzles and clean separately. Rinse inside the tank thoroughly using a pressure hose and flush system with clean water.

RESISTANCE WARNING

GROUP	F I	HERBICIDE
-------	------------	-----------

Spearhead Selective Herbicide is a member of the nicotinilide, phenoxy and pyridine groups of herbicides. Spearhead Selective Herbicide has the inhibitors of carotenoid biosynthesis and disruptor of cell growth modes of action. For weed resistance management Spearhead is a group **F** and **I** herbicide. Some naturally occurring weed biotypes resistant to Spearhead and other Group **F** and **I** herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Spearhead or other Group **F** or **I** herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Bayer CropScience accepts no liability for any losses that may result from the failure of Spearhead Selective Herbicide to control resistant weeds.

COMPATIBILITY

No information is available that indicates physical compatibility with other products. Tank-mixing other products with Spearhead cannot therefore be recommended. As formulations of other manufacturers' products are beyond the control of Bayer CropScience, all mixtures should be tested prior to mixing commercial quantities.

PRECAUTION

Re-Entry Period

Do not allow entry into treated areas until the spray has dried. When prior entry is necessary wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

Manual handling of turf

Hand-weeding and transplanting should not be performed before 2 weeks after spray application, unless workers wear overalls (or equivalent clothing) and chemical resistant gloves.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Spray drift from Spearhead application may cause injury to herbaceous plants, vegetables, fruit crops, vines, ornamentals and trees.

DO NOT use clippings from treated areas for mulch around vegetables or fruit trees.

PROTECTION OF LIVESTOCK

DO NOT graze treated turf or feed turf clippings from any treated area to poultry or livestock.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish and other aquatic life. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant local, State or Territory government regulations. DO NOT burn empty containers or product. DO NOT re-use container for any other purpose.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage eyes. Will irritate skin. Avoid contact with eyes and skin. If product in eyes, wash it out immediately with water. Wash hands after use. When opening the container, preparing the spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves and goggles. After each days use wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13 11 26.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet.

Exclusion of Liability

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

APVMA Approval No.: 53833 / 60388

Spearhead is a Registered Trademark of the Bayer Group



**FOR 24 HOUR SPECIALIST ADVICE
IN EMERGENCY ONLY
PHONE 1800 033 111**

Bayer CropScience Pty Ltd
A.B.N. 87 000 226 022
391-393 Tooronga Rd
Hawthorn East, Vic. 3123
Telephone: (03) 9248 6888 Technical Enquiries: 1800 804 479
Website: www.environmentalscience.bayer.com.au

BATCH NO:

DATE OF MANUFACTURE: